



Canadian Water Quality Guidelines for the Protection of Aquatic Life

SIMAZINE

Simazine ($C_7H_{12}ClN_5$) is a triazine herbicide with a CAS name and number of 2-chloro-4,6-bis(ethylamino)-1,3,5-triazine and 122-34-9, respectively. Simazine was first registered in Canada in 1963 under the trade names Simadex, Simmaprim, and Princep (Agriculture and Agri-Food Canada 1997).

Simazine is a selective herbicide used for the control of annual broadleaf and grass weeds in raspberries, loganberries, highbush blueberries, apples, pears, grapes, asparagus, and ornamentals. Noncrop uses include total weed control in industrial areas, at airports, and along shelterbelts and rights-of-way, and aquatic weed control in ditches, farm ponds, fish hatcheries, aquaria, and fountains (Agriculture and Agri-Food Canada 1997). Its mode of action is to inhibit photosynthetic electron transport (Tomlin 1994).

In 1987, 219 t of formulated simazine and 1684 t of technical grade simazine were imported into Canada (Statistics Canada 1988). With the exception of spills, concentrations of simazine detected in Canadian freshwater sources range from 0.0003 to $6 \mu\text{g}\cdot\text{L}^{-1}$ (Frank et al. 1979; Frank 1981). Simazine contamination of freshwater may result from spraying directly into watercourses, vapour drift, and precipitation, or from surface runoff and groundwater intrusions from treated lands. Accidents and spills of simazine may also contaminate surface waters (CCME 1991a).

The major fate processes of simazine are microbial degradation and, possibly, photodegradation combined with sorption to sediments and aquatic plants (Muir 1991). Volatilization to the atmosphere is not a major fate process (Henry's law constant of $0.00034 \text{ Pa}\cdot\text{m}^3\cdot\text{mol}^{-1}$) (Suntio et al. 1988). The hydrolysis half-life of simazine is 70 d at pH 5 and >200 d at pH 7 and 9 (Burkhard and Guth 1981). Simazine can be relatively persistent in aquatic systems, particularly shallow, well-mixed lakes and ponds (Jenkins and Buikema 1990).

Simazine does not bioaccumulate or biomagnify in the food web. The bioaccumulation potential is low as evidenced by BCFs <100. The depuration half-life in fish is <7 d following exposure if the organism is transferred to uncontaminated water, indicating that simazine is

rapidly excreted or metabolized (Rodgers 1970; Mayer and Sanders 1977; Niimi 1987).

Water Quality Guideline Derivation

The Canadian water quality guideline for simazine for the protection of freshwater life was developed based on the CCME protocol (CCME 1991b).

Freshwater Life

Simazine has a low toxicity to fish (WSSA 1983). The USDA (1984) concluded that simazine should not affect fish at concentrations below its water solubility. Reported 48-h LC_{50} s range from $>10 \text{ mg}\cdot\text{L}^{-1}$ for medaka (Hashimoto and Nishiuchi 1981) to $85 \text{ mg}\cdot\text{L}^{-1}$ for rainbow trout (Alabaster 1969). Snow (1963) reported that simazine was not toxic to zooplankton and other animal life composing the diet of fish being cultured in the ponds. Laboratory tests on bottom organisms gave an acute LD_{50} of $28 \text{ mg}\cdot\text{L}^{-1}$ (Walker 1964).

Marchini et al. (1988) found the acute toxicity of simazine to *Daphnia magna* to be $>3.5 \text{ mg}\cdot\text{L}^{-1}$ using 24-h and 48-h immobilization tests. Sanders (1970) found that *D. magna* and seed shrimp (*Cypridopsis vidua*) were immobilized after 48-h exposures to simazine concentrations of 1.0 and $3.2 \text{ mg}\cdot\text{L}^{-1}$, respectively.

The USDA's (1984) review of simazine lists LC_{50} s for stonefly larvae (*Pteronarcys* sp.) of 1.9 (96-h) to 50 (48-h) $\text{mg}\cdot\text{L}^{-1}$ and a 24-h LC_{50} for the freshwater copepod *Heliodiaptomus viduus* of $1.0 \text{ mg}\cdot\text{L}^{-1}$. While conducting 48-h LC_{50} tests with *D. pulex*, Fitzmayer et al.

Table 1. Water quality guidelines for simazine for the protection of aquatic life (CCME 1991a).

| Aquatic life | Guideline value ($\mu\text{g}\cdot\text{L}^{-1}$) |
|--------------|---|
| Freshwater | 10 |
| Marine | NRG* |

*No recommended guideline.




Figure 1. Select freshwater toxicity data for simazine.

(1982) noted that the daphnids became sedentary at simazine concentrations of 1 to 50 $\text{mg}\cdot\text{L}^{-1}$. For *D. pulex* and *Moina macrocopa*, 3-h LC₅₀s were >40 $\text{mg}\cdot\text{L}^{-1}$ (Hashimoto and Nishiuchi 1981). A single amphibian toxicity (48-h TL_m) was >100 $\text{mg}\cdot\text{L}^{-1}$ for the tadpole of *Bufo bufo japonicus* (Hashimoto and Nishiuchi 1981).

Chronic NOELs for *Daphnia* sp. and mud crabs were 4 $\text{mg}\cdot\text{L}^{-1}$ and 1000 $\text{mg}\cdot\text{L}^{-1}$, respectively (USDA 1984).

Kosanke et al. (1988) examined the effect of simazine on the ontogenesis of freshwater snail (*Lymnaea stagnalis*) embryos. At 20.1 $\mu\text{g}\cdot\text{L}^{-1}$ simazine, all eggs were killed (762 eggs). In other tests with molluscs, a 48-h LC₅₀ value >100 $\text{mg}\cdot\text{L}^{-1}$ for the snails *Indoplanorbis exustus*, *Semisulcospira libertina*, and *Physa acuta* were reported (Hashimoto and Nishiuchi 1981).

After extensive studies of in situ enclosures, Goldsborough and Robinson (1986), arrived at a minimum periphyton community LC₅₀ of 100 $\mu\text{g}\cdot\text{L}^{-1}$. Data indicated rapid recovery of the organisms within the enclosures even after treatment with 5.0 $\text{mg}\cdot\text{L}^{-1}$. Therefore, even though simazine may exert adverse effects on the organisms that form the basis of the aquatic food web, these effects are transient and do not translate to adverse effects on the organisms depending on the plankton community for food. The study by Kosanke et al. (1988), where effects of simazine were reported at

concentrations of 20.1 $\mu\text{g}\cdot\text{L}^{-1}$, was not used to establish the water quality guideline because no response curve was available for that study. The water quality guideline for simazine for the protection of freshwater life is 10 $\mu\text{g}\cdot\text{L}^{-1}$. It was derived by multiplying the minimum plankton community LC₅₀ of 100 $\mu\text{g}\cdot\text{L}^{-1}$ (Goldsborough and Robinson 1986) by a safety factor of 0.1 (CCME 1991a).

References

- Agriculture and Agri-Food Canada. 1997. Regulatory Information on Pesticide Products (RIPP) Database (CCINFODISK). Produced by Agriculture and Agri-Food Canada and distributed by the Canadian Centre for Occupational Health and Safety. CD-ROM.
- Alabaster, J.S. 1969. Survival of fish in 164 herbicides, insecticides, fungicides, wetting agents and miscellaneous substances. *Int. Pest Control*. 11(2):29–35.
- Burkhard, N., and J.A. Guth. 1981. Chemical hydrolysis of 2-chloro-4, 6-bis-(alkylamino)-1,3,5-triazine herbicides and their breakdown in soil under the influence of adsorption. *Pestic. Sci.* 12:45–52.
- CCME (Canadian Council of Ministers of the Environment). 1991a. Appendix VIII—Canadian water quality guidelines: Updates (April 1991), metolachlor, simazine, and captan. In: Canadian water quality guidelines, Canadian Council of Resource and Environment Ministers. 1987. Prepared by the Task Force on Water Quality Guidelines.
- . 1991b. Appendix IX—A protocol for the derivation of water quality guidelines for the protection of aquatic life (April 1991). In: Canadian water quality guidelines, Canadian Council of Resource and Environment Ministers. 1987. Prepared by the Task Force on Water Quality Guidelines. [Updated and reprinted with minor revisions and editorial changes in Canadian environmental quality guidelines, Chapter 4, Canadian Council of Ministers of the Environment, 1999, Winnipeg.]
- Fitzmayer, K.M., J.G. Geiger, and M.J. van den Avyle. 1982. Acute toxicity effects of simazine on *Daphnia pulex* and larval striped bass. In: *Proc. Annu. Conf. Southeast. Assoc. Fish and Wild. Agencies*, 1982:146–156.
- Frank, R. 1981. Pesticides and PCB in the Grand and Saugeen river basins. *J. Gt. Lakes Res.* 7(4):440–454.
- Frank, R., G.J. Sirons, R.L. Thomas, and K. McMillan. 1979. Triazine residues in suspended solids (1974–1976) and water (1977) from the mouths of Canadian streams flowing into the Great Lakes. *J. Gt. Lakes Res.* 5(2):131–138.
- Goldsborough, L.G., and G.G.C. Robinson. 1986. Changes in periphytic algal community structure as a consequence of short herbicide exposures. *Hydrobiologia* 139:177–192.
- Hashimoto, Y., and Y. Nishiuchi. 1981. Establishment of bioassay methods for the evaluation of acute toxicity of pesticides to aquatic organisms. *J. Pestic. Sci.* 6:257–264. (In Japanese with English summary.)
- Jenkins, D.G., and A.L.J. Buikema. 1990. Response of a winter plankton food web to simazine. *Environ. Toxicol. Chem.* 9(6):693–705.
- Kosanke, G.J., W.W. Schwippert, and T.W. Beneke. 1988. The impairment of mobility and development in freshwater snails (*Physa fontinalis* and *Lymnaea stagnalis*) caused by herbicides. *Comp. Biochem. Physiol.* 90C(2):373–379.
- Marchini, S., L. Passerini, D. Cesareo, and M.L. Tosato. 1988. Herbicidal triazines: Acute toxicity on *Daphnia*, fish, and plants and

- analysis of its relationships with structural factors. *Ecotoxicol. Environ. Saf.* 16:148–157.
- Mayer, F.L., and H.O. Sanders. 1977. Simazine effects on non-target aquatic organisms: A preliminary report. *Proc. South. Weed Sci. Soc.* 30:333 (abstract).
- Muir, D.C.G. 1991. Dissipation and transformations in water and sediments In: *Environmental chemistry of herbicides*, Vol. 2, R. Grover, ed. CRC Press, Boca Raton, FL.
- Niimi, A.J. 1987. Biological half-lives of chemicals in fishes. *Rev. Environ. Contam. Toxicol.* 99:1–46.
- Rodgers, C.A. 1970. Uptake and elimination of simazine by green sunfish (*Lepomis cyanellus* Raf.). *Weed Sci.* 18:134–136.
- Sanders, H.O. 1970. Toxicities of some herbicides to six species of freshwater crustaceans. *J. Water Pollut. Cont. Fed.* 42(8):1544–1550.
- Snow, J.R. 1963. Simazine as an algicide for bass ponds. *Prog. Fish-Cult.* 25(1):34–36.
- Statistics Canada. 1988. Imports: Commodity by country. C.I.T.C. detail 1987–1988. Statistics Canada, International Trade Division, Ottawa.
- Suntio, L.R., W.Y. Shiu, D. Mackay, J.N. Seiber and D. Glotfelty. 1988. Critical review of Henry's law constants for pesticides. *Rev. Environ. Contam. Toxicol.* 109:1–59.
- Tomlin, C. (ed.). 1994. *The pesticide manual: A world compendium*. 10th ed. (Incorporating the *Agrochemicals handbook*.) British Crop Protection Council and Royal Society of Chemistry, Thornton Heath, UK.
- USDA (U.S. Department of Agriculture). 1984. Simazine. In: *Pesticide background statements, Agricultural handbook No. 633, Vol. 1, Herbicides*. USDA, Forest Service, Washington, DC.
- Walker, C.R. 1964. Simazine and other s-triazine compounds as aquatic herbicides in fish habitats. *Weeds* 12(2):134–149.
- WSSA (Weed Science Society of America.). 1983. *Herbicide handbook*. 5th ed. WSSA, Champaign, IL.

Reference listing:

Canadian Council of Ministers of the Environment. 1999. Canadian water quality guidelines for the protection of aquatic life: Simazine. In: *Canadian environmental quality guidelines, 1999*, Canadian Council of Ministers of the Environment, Winnipeg.

For further scientific information, contact:

Environment Canada
Guidelines and Standards Division
351 St. Joseph Blvd.
Hull, QC K1A 0H3
Phone: (819) 953-1550
Facsimile: (819) 953-0461
E-mail: ceqg-rcqe@ec.gc.ca
Internet: <http://www.ec.gc.ca>

For additional copies, contact:

CCME Documents
c/o Manitoba Statutory Publications
200 Vaughan St.
Winnipeg, MB R3C 1T5
Phone: (204) 945-4664
Facsimile: (204) 945-7172
E-mail: spcme@chc.gov.mb.ca